

Spring 2014 Math 272 Exam 2 Review Sheet

The exam will be in class on Wednesday, April 2, and will cover Sections 3.1-4.4.

You will not be allowed use of a calculator or any other device other than your pencil or pen and some scratch paper. Notes are also not allowed. In kindness to your fellow test-takers, please turn off all cell phones and anything else that might beep or be a distraction.

Important topics:

- Determinants
- Eigenvalues and eigenvectors
- Application to matrix powers
- Vector spaces and subspaces (of vectors, matrices, and polynomials)
- Span
- Linear independence
- Basis

If you have questions on certain topics or want more practice on exam-like problems, please feel free to stop by my office.

Chapter 3 review (p227): 4, 7a, 8a-f, 10, 12, 18, 19, 20

Chapter 4 review (p316): 3, 5, 6, 8, 9, 13, 15, 16, 20, 21, 22, 27

Extra review problems:

1. Prove that $|\mathbf{A}^{-1}|=1/|\mathbf{A}|$ for all invertible matrices \mathbf{A} .
2. Use induction to prove that if λ is an eigenvalue of a matrix \mathbf{A} with corresponding eigenvector \mathbf{v} , then λ^m is an eigenvalue of \mathbf{A}^m with corresponding eigenvector \mathbf{v} .
3. Prove that if \mathbf{A} is nilpotent ($\mathbf{A}^m=0$ for some positive integer m), then 0 is an eigenvalue of \mathbf{A} . In fact, show that 0 is the *only* eigenvalue of \mathbf{A} .

4. Let \mathbf{A} be a 2x2 matrix with eigensystem $\left\{ \left\{ \lambda_1 = -\frac{1}{2}, \mathbf{v}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}, \left\{ \lambda_2 = 1, \mathbf{v}_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\} \right\}$
and let $\mathbf{x}_0 = \begin{bmatrix} 3 \\ 3 \end{bmatrix}$. Find the limit of $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0$ as $k \rightarrow \infty$.

5. Suppose a 2×2 matrix \mathbf{A} has eigensystem $\left\{ \left\{ \lambda_1 = 1, \mathbf{v}_1 = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \right\}, \left\{ \lambda_2 = \frac{2}{3}, \mathbf{v}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\} \right\}$.

Determine $\lim_{k \rightarrow \infty} \mathbf{A}^k \mathbf{x}_0$ if $\mathbf{x}_0 = \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix}$.

6. Find a basis for the vector space $\{p(x) = a_0 + a_1x + a_2x^2 : p(1) = 0\}$. What is the dimension of this space?
7. Find a basis for the space $\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a + b - c = 0 \text{ and } b + 2d = 0 \right\}$. What is the dimension of this space?
8. Prove that $B = \{1 + x, x + x^2, 1 - x\}$ is a basis for \mathbf{P}_2 . Express $2 + x + x^2$ as a linear combination of the vectors in this basis.
9. Determine if the set S is a subspace of the vector space V . If it is, then find a basis for S . If it is not, then explain why not.

(a) $S = \left\{ \begin{pmatrix} a & b \\ a & b \end{pmatrix} : a, b \text{ are real} \right\}$ for $V =$ set of all 2×2 matrices with real entries

(b) $S = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} : x + z = 1 \right\}$ for $V = \mathbf{R}^3$

Partial solutions

- Let \mathbf{A} be an invertible matrix, so $\mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$. Then $1 = |\mathbf{I}| = |\mathbf{A}\mathbf{A}^{-1}| = |\mathbf{A}| |\mathbf{A}^{-1}|$, so $|\mathbf{A}^{-1}| = 1/|\mathbf{A}|$. Note that we can divide by $|\mathbf{A}|$ since invertible matrices are nonsingular ($|\mathbf{A}| \neq 0$).
- Suppose $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$. We will prove that $\mathbf{A}^m\mathbf{v} = \lambda^m\mathbf{v}$ for any positive integer m by induction. The base case is $m=1$, and we already know $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$. Suppose $\mathbf{A}^m\mathbf{v} = \lambda^m\mathbf{v}$ for some positive integer m . Multiply both sides of this equation by \mathbf{A} to obtain $\mathbf{A}^{m+1}\mathbf{v} = \lambda^m\mathbf{A}\mathbf{v} = \lambda^m\lambda\mathbf{v} = \lambda^{m+1}\mathbf{v}$. Hence the statement is also true for $m+1$. Therefore, by induction the statement is true for all positive integers m .

- Suppose $\mathbf{A}^m = \mathbf{0}$ for some positive integer m . Then $|\mathbf{A}|^m = |\mathbf{A}^m| = 0$, so $|\mathbf{A} - 0\mathbf{I}| = |\mathbf{A}| = 0$. This says that $\lambda = 0$ satisfies the characteristic equation of \mathbf{A} , so is an eigenvalue of \mathbf{A} . Now suppose that λ is any eigenvalue of \mathbf{A} . For some nonzero vector \mathbf{v} , $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$. From #2, $\mathbf{A}^m\mathbf{v} = \lambda^m\mathbf{v}$. But $\mathbf{A}^m = \mathbf{0}$, so $\lambda^m\mathbf{v} = \mathbf{A}^m\mathbf{v} = \mathbf{0}\mathbf{v} = \mathbf{0}$. We know \mathbf{v} is nonzero, so it must be that $\lambda = 0$. Hence $\lambda = 0$ is the only eigenvalue of \mathbf{A} .
- Write \mathbf{x}_0 as a linear combination of eigenvectors then multiply by \mathbf{A}^k and simplify using eigenvalues. The limit is $\begin{bmatrix} 4 \\ 2 \end{bmatrix}$.
- The limit is $\begin{bmatrix} 3/4 \\ 1/4 \end{bmatrix}$.
- A basis is $B = \{x-1, x^2-1\}$, so dimension is 2.
- A basis is $B = \left\{ \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 2 & -2 \\ 0 & 1 \end{bmatrix} \right\}$, so dimension is 2.
- To prove the set is a basis, we first note that it contains 3 vectors and \mathbf{P}_2 is a 3 dimensional space, so we only need to prove it spans \mathbf{P}_2 or that it is linearly independent. To do the latter, show that the only linear combination of the 3 polynomials that equals zero has all 3 coefficients equal to zero. $2+x+x^2$ equals the sum of the 3 basis polynomials.
- (a) Yes, this is a subspace; (b) Not a subspace since doesn't contain zero vector.

Please email me if you find any errors in the solutions. Thank you!

Practice exam:

- Compute $\begin{vmatrix} 1 & 0 & 0 \\ p & 1 & s \\ q & r & 2 \end{vmatrix}$. For what values of p, q, r , and s is the following set a basis for \mathbf{R}^3 : $\left\{ \begin{bmatrix} 1 \\ p \\ q \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ r \end{bmatrix}, \begin{bmatrix} 0 \\ s \\ 2 \end{bmatrix} \right\}$

- Prove that $B = \{1-x, 1+x, x-x^2\}$ is a basis for \mathbf{P}_2 . Express $3x^2+x$ as a linear combination of the vectors in this basis.

- Find a basis for the space $\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a+3b=0 \text{ and } c+3d=0 \right\}$. What is the dimension of this space?

- Suppose V is a vector space and $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is a basis for V . Prove that $\{\mathbf{u}, \mathbf{u}+\mathbf{v}, \mathbf{u}+\mathbf{v}+\mathbf{w}\}$ is also a basis for V .

- Prove that $\lambda=0$ is an eigenvalue of a matrix \mathbf{A} if and only if \mathbf{A} is singular.