

A nilpotent Whitehead theorem for TQ-homology of structured ring spectra

Michael Ching¹, John E. Harper²

¹ Department of Mathematics and Statistics, Amherst College, Amherst, MA, 01002, USA

² Department of Mathematics, The Ohio State University, Newark, 1179 University Dr, Newark, OH 43055, USA
E-mail: mching@amherst.edu, harper.903@math.osu.edu

Abstract

The aim of this short paper is to prove a TQ-Whitehead theorem for nilpotent structured ring spectra. We work in the framework of symmetric spectra and algebras over operads in modules over a commutative ring spectrum. Our main result can be thought of as a TQ-homology analog for structured ring spectra of Dror’s generalized Whitehead theorem for topological spaces; here TQ-homology is short for topological Quillen homology. We also prove retract theorems for the TQ-completion and homotopy completion of nilpotent structured ring spectra.

2010 Mathematics Subject Classification. **55P43**. 55U35, 55P48, 18G55

Keywords. Symmetric spectra, structured ring spectra, calculus of functors, operads, topological Quillen homology.

1 Introduction

In this paper we work with algebraic structures in symmetric spectra, or \mathcal{R} -modules, that can be described as algebras over an operad \mathcal{O} which is reduced, meaning that $\mathcal{O}[0] = *$, the trivial \mathcal{R} -module; such \mathcal{O} -algebras are non-unital. Here, we assume that \mathcal{R} is any commutative monoid object in the category $(\mathbf{Sp}^{\Sigma}, \otimes_S, S)$ of symmetric spectra [19, 31]; in other words, \mathcal{R} is a commutative ring spectrum, and we denote by $(\mathbf{Mod}_{\mathcal{R}}, \wedge, \mathcal{R})$ the closed symmetric monoidal category of \mathcal{R} -modules.

Topological Quillen homology, or TQ-homology for short, is the analog for \mathcal{O} -algebras of the ordinary integral homology of spaces. There is an extensive literature, but it is useful to start with [13, 25, 29], followed by [11, 12, 16, 30] along with [2, 3, 4, 20, 22, 23], among others. A study of the homotopy completion tower, which is weakly equivalent to the Goodwillie Taylor tower [14, 21] of the identity functor on \mathcal{O} -algebras (see, for instance, [18, 20, 24, 27]), is carried out in [18], where the TQ-completion construction is introduced as the analog for \mathcal{O} -algebras of the Bousfield-Kan [5] integral completion of spaces; a study of TQ-completion is carried out in [6, 7] for connected \mathcal{O} -algebras. The homotopy completion tower is exploited in [18] to establish a TQ-Serre Finiteness Theorem; also proved, using similar tools, are a TQ-Hurewicz theorem and a TQ-Whitehead theorem, which generalize earlier results in this direction to \mathcal{O} -algebras. These are analogs of the Serre Finiteness, Hurewicz, and Whitehead theorems that are well-known for spaces; each of these theorems for TQ-homology starts with the assumption that the \mathcal{O} -algebra is connected.

In this paper we explore the possibility of removing the connectivity assumption from an \mathcal{O} -algebra, in the case of the TQ-Whitehead theorem, by replacing “connected \mathcal{O} -algebra” with “nilpotent \mathcal{O} -algebra”; this is motivated by Dror’s generalized Whitehead theorem for spaces [8] which replaced “simply connected space” with “nilpotent space” in the original Whitehead theorem for spaces.

Tbilisi Mathematical Journal 11(3) (2018), pp. 69–79.
Tbilisi Centre for Mathematical Sciences.

Received by the editors: 26 June 2018.
Accepted for publication: 11 July 2018.

Our approach, which is somewhat indirect and was motivated by the work in [1], is to play off the homotopy completion and TQ-completion constructions against each other: more precisely, we resolve the cosimplicial TQ-resolution of an \mathcal{O} -algebra X , whose homotopy limit is the TQ-completion of X , term-by-term via the homotopy completion tower, but with respect to the operad $\tau_{M-1}\mathcal{O}$ characterizing M -nilpotent \mathcal{O} -algebras (Remark 2.5). We use this to establish a nilpotent TQ-completion retract result (Theorem 2.12), and as a consequence, we establish the following nilpotent TQ-Whitehead theorem.

Theorem 1.1 (Nilpotent TQ-Whitehead theorem). Let \mathcal{O} be an operad in \mathcal{R} -modules with $\mathcal{O}[0] = *$. Let $M \geq 2$ and suppose X, Y are M -nilpotent \mathcal{O} -algebras (Definition 2.3). Consider any map $f: X \rightarrow Y$ of \mathcal{O} -algebras. Then f is a weak equivalence if and only if f induces a weak equivalence $\mathrm{TQ}(X) \simeq \mathrm{TQ}(Y)$ on TQ-homology. Here we are assuming that \mathcal{O} satisfies Assumption 2.1.

To keep this paper appropriately concise, we freely use notation from [18].

Acknowledgments

The second author would like to thank Greg Arone, Bill Dwyer, and Emmanuel Farjoun for helpful remarks at an early stage of this project. The authors would like to thank Andy Baker and Birgit Richter for the opportunity to announce these results at the 2011 Conference on Structured Ring Spectra in Hamburg, and an anonymous referee for helpful comments and suggestions. The first author was supported in part by National Science Foundation grant DMS-1308933.

2 Nilpotent structured ring spectra

The following basic assumption ensures, in particular, that the forgetful functor from \mathcal{O} -algebras to the underlying category of \mathcal{R} -modules preserves cofibrant objects; see, for instance, [26, 28, 33].

Assumption 2.1. Let \mathcal{O} be an operad in \mathcal{R} -modules and consider the associated unit map $I \rightarrow \mathcal{O}$. Assume that $I[r] \rightarrow \mathcal{O}[r]$ is a flat stable cofibration between flat stable cofibrant objects in $\mathrm{Mod}_{\mathcal{R}}$ for each $r \geq 0$ ([18, 7.7]).

Recall from [18] that associated to the operad \mathcal{O} is the tower of operads

$$\tau_1\mathcal{O} \leftarrow \tau_2\mathcal{O} \leftarrow \cdots \leftarrow \tau_{n-1}\mathcal{O} \leftarrow \tau_n\mathcal{O} \leftarrow \cdots \quad (2.1)$$

where

$$(\tau_n\mathcal{O})[t] := \begin{cases} \mathcal{O}[t], & \text{for } t \leq n, \\ *, & \text{otherwise,} \end{cases}$$

In other words, $\tau_n\mathcal{O}$ is the operad built from \mathcal{O} by truncating \mathcal{O} above level n , where $*$ denotes the trivial \mathcal{R} -module. If X is an \mathcal{O} -algebra, then the tower (2.1) of operads induces the *completion tower* [18, 3.8] of X

$$\tau_1\mathcal{O} \circ_{\mathcal{O}} (X) \leftarrow \tau_2\mathcal{O} \circ_{\mathcal{O}} (X) \leftarrow \cdots \leftarrow \tau_{n-1}\mathcal{O} \circ_{\mathcal{O}} (X) \leftarrow \tau_n\mathcal{O} \circ_{\mathcal{O}} (X) \leftarrow \cdots \quad (2.2)$$

of \mathcal{O} -algebras whose limit is the *completion* X^\wedge of X . To make this construction homotopy meaningful, one can derive everything in sight by constructing the *homotopy completion tower* [18, 3.13] of X

$$\tau_1\mathcal{O} \circ_{\mathcal{O}}^h (X) \leftarrow \tau_2\mathcal{O} \circ_{\mathcal{O}}^h (X) \leftarrow \cdots \leftarrow \tau_{n-1}\mathcal{O} \circ_{\mathcal{O}}^h (X) \leftarrow \tau_n\mathcal{O} \circ_{\mathcal{O}}^h (X) \leftarrow \cdots \quad (2.3)$$

of \mathcal{O} -algebras whose homotopy limit is the *homotopy completion* X^{h^\wedge} of X .

Remark 2.2. For instance, if \mathcal{O} is the operad whose algebras are the non-unital commutative algebra spectra (i.e., where $\mathcal{O}[t] = \mathcal{R}$ for each $t \geq 1$ and $\mathcal{O}[0] = *$), then the tower (2.2) is isomorphic to the usual X -adic completion of X tower of the form

$$X/X^2 \leftarrow X/X^3 \leftarrow \cdots \leftarrow X/X^n \leftarrow X/X^{n+1} \leftarrow \cdots \quad (2.4)$$

where X/X^n denotes the \mathcal{O} -algebra with underlying \mathcal{R} -module defined by the pushout diagram ($n \geq 2$)

$$\begin{array}{ccc} X^{\wedge n} & \xrightarrow{m_n} & X \\ \downarrow & & \downarrow \\ * & \longrightarrow & X/X^n \end{array}$$

in \mathcal{R} -modules; here, m_n denotes the usual multiplication map.

Definition 2.3. Let X be an \mathcal{O} -algebra and $M \geq 2$. We say that X is M -nilpotent if the map of t -ary operations $\mathcal{O}[t] \wedge X^{\wedge t} \rightarrow X$ factors as

$$\mathcal{O}[t] \wedge X^{\wedge t} \rightarrow * \rightarrow X \quad \text{for each } t \geq M,$$

where $*$ denotes the trivial \mathcal{R} -module; in other words, if all the M -ary operations of X and higher are trivial.

Remark 2.4. Each term X/X^n in (2.4) is an n -nilpotent non-unital commutative algebra spectrum, and similarly, each term $\tau_n \mathcal{O} \circ_{\mathcal{O}} (X)$ in the completion tower (2.2) is an $(n + 1)$ -nilpotent \mathcal{O} -algebra.

Remark 2.5. Since M -nilpotent \mathcal{O} -algebras ($M \geq 2$) are the same as $\tau_{M-1} \mathcal{O}$ -algebras equipped with \mathcal{O} -action maps (Proposition 2.6), for ease of notational purposes (in the various subscripts) we often state assumptions in terms of $(N + 1)$ -nilpotent \mathcal{O} -algebras ($N \geq 1$).

Proposition 2.6. If X is an \mathcal{O} -algebra and $N \geq 1$, then X is $(N + 1)$ -nilpotent if and only if the left \mathcal{O} -action map $\mathcal{O} \circ (X) \rightarrow X$ factors as

$$\mathcal{O} \circ (X) \rightarrow (\tau_N \mathcal{O}) \circ (X) \rightarrow X$$

through a $\tau_N \mathcal{O}$ -algebra structure on X . In other words, $(N + 1)$ -nilpotent \mathcal{O} -algebras are the same as $\tau_N \mathcal{O}$ -algebras equipped with \mathcal{O} -action maps induced by restriction along $\mathcal{O} \rightarrow \tau_N \mathcal{O}$.

Proof. This follows immediately from Definition 2.3. Q.E.D.

Remark 2.7. It is natural to ask, if we start with a small diagram of $\tau_N \mathcal{O}$ -algebras, and we compute its limit (resp. colimit) in \mathcal{O} -algebras versus in $\tau_N \mathcal{O}$ -algebras, when are they isomorphic? Since all small limits, filtered colimits, and reflexive coequalizers are created in the underlying category of \mathcal{R} -modules by the forgetful functor (see, for instance, [15, 17]), for both \mathcal{O} -algebras and $\tau_N \mathcal{O}$ -algebras, it follows that $(N + 1)$ nilpotent \mathcal{O} -algebras are closed under all small limits, filtered colimits, and reflexive coequalizers. On the other hand, since cofibrant replacement of an $(N + 1)$ -nilpotent \mathcal{O} -algebra, as an \mathcal{O} -algebra, involves gluing on cells of \mathcal{O} -algebras that are not nilpotent,

in general, a left derived functor evaluated on a fixed $(N + 1)$ -nilpotent \mathcal{O} -algebra will often take on different values depending on whether it is left derived on \mathcal{O} -algebras versus on $\tau_N \mathcal{O}$ -algebras. Note that since the category of $\tau_1 \mathcal{O}$ -algebras is isomorphic to the category of left $\mathcal{O}[1]$ -modules [18], 2-nilpotent \mathcal{O} -algebras are the same as left $\mathcal{O}[1]$ -modules equipped with a trivial \mathcal{O} -algebra structure induced by restriction along $\mathcal{O} \rightarrow \tau_1 \mathcal{O}$.

If $N \geq 1$, repeated application of the functorial factorization construction in [18, 5.48] shows that the upper horizontal finite tower in (2.5) may be factored as

$$\begin{array}{ccccccccc} \mathcal{O} & \longrightarrow & \tau_N \mathcal{O} & \longrightarrow & \tau_{N-1} \mathcal{O} & \longrightarrow & \cdots & \longrightarrow & \tau_2 \mathcal{O} & \longrightarrow & \tau_1 \mathcal{O} & (2.5) \\ & \searrow^{(*)} & \uparrow^{(\#)} \simeq & & \uparrow^{(\#)} \simeq & & & & \uparrow^{(\#)} \simeq & & \uparrow^{(\#)} \simeq & \\ & & J_N & \longrightarrow & J_{N-1} & \longrightarrow & \cdots & \longrightarrow & J_2 & \longrightarrow & J_1 \end{array}$$

where the maps $(*)$ are cofibrations of operads, the maps $(\#)$ are weak equivalences of operads, and J_n is reduced for each $1 \leq n \leq N$.

The following is proved in [18, 5.49(b)].

Proposition 2.8. Let $1 \leq n < N$. If $X \in \mathbf{Alg}_{J_N}$ is cofibrant, then X is also cofibrant in $\mathbf{Alg}_{\mathcal{O}}$. Similarly, if $X \in \mathbf{Alg}_{J_n}$ is cofibrant, then X is also cofibrant in $\mathbf{Alg}_{J_{n+1}}$.

If X is an $(N+1)$ -nilpotent \mathcal{O} -algebra, then its \mathcal{O} -algebra structure is induced from its underlying $\tau_N \mathcal{O}$ -algebra structure via the composite map

$$\mathcal{O} \circ (X) \rightarrow \tau_N \mathcal{O} \circ (X) \rightarrow X.$$

Hence we may start with any $(N + 1)$ -nilpotent \mathcal{O} -algebra X , and consider the functorial cofibrant replacement

$$* \rightarrow \tilde{X} \xrightarrow{\simeq} X \quad (2.6)$$

of X in \mathbf{Alg}_{J_N} , which is also a cofibrant replacement of X in $\mathbf{Alg}_{\mathcal{O}}$; we are using the positive flat stable model structure (see, for instance, [15, 18]).

Definition 2.9. If X is an $(N + 1)$ -nilpotent \mathcal{O} -algebra, then its TQ -homology (resp. $\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}$ -homology) is the \mathcal{O} -algebra

$$\begin{aligned} \mathrm{TQ}(X) &:= \tau_1 \mathcal{O} \circ_{\mathcal{O}}^h (X) \simeq J_1 \circ_{\mathcal{O}} (\tilde{X}) \\ \text{resp. } \mathrm{TQ}|_{\mathrm{Nil}_{N+1}}(X) &:= \tau_1 \mathcal{O} \circ_{\tau_N \mathcal{O}}^h (X) \simeq J_1 \circ_{J_N} (\tilde{X}) \end{aligned}$$

Here, “ TQ -homology” is an abbreviation for “topological Quillen homology”.

Remark 2.10. In other words, $\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}(X)$ is the topological Quillen homology of X (i.e., the derived indecomposable quotient of X) with respect to $\tau_N \mathcal{O}$ -algebras, while $\mathrm{TQ}(X)$ is the the derived indecomposable quotient of X with respect to \mathcal{O} -algebras; it may be helpful to note that $\tau_1 \tau_N \mathcal{O} = \tau_1 \mathcal{O}$. The indicated weak equivalences are consequences of [18, 4.10].

This construction comes with a TQ-Hurewicz map of the form $X \rightarrow \mathrm{TQ}(X)$, given by the map $\tilde{X} \rightarrow J_1 \circ_{\mathcal{O}} (\tilde{X})$ in $\mathrm{Alg}_{\mathcal{O}}$, and iterating this Hurewicz map leads to a cosimplicial resolution of X with respect to TQ-homology of the form

$$X \longrightarrow \mathrm{TQ}(X) \rightrightarrows (\mathrm{TQ})^2(X) \rightrightarrows (\mathrm{TQ})^3(X) \cdots$$

(see Remark 2.11) given by the map of cosimplicial \mathcal{O} -algebras

$$\tilde{X} \longrightarrow J_1 \circ_{\mathcal{O}} (\tilde{X}) \rightrightarrows J_1 \circ_{\mathcal{O}} J_1 \circ_{\mathcal{O}} (\tilde{X}) \rightrightarrows \cdots \quad (2.7)$$

where \tilde{X} is regarded as a constant cosimplicial diagram. Recall [6, 18] that applying $\mathrm{holim}_{\Delta}(-)$ to the map of cosimplicial \mathcal{O} -algebras in (2.7) gives the TQ-completion map of the form $X \rightarrow X_{\mathrm{TQ}}^{\wedge}$.

Remark 2.11. Throughout this paper we have suppressed the codegeneracy maps from the diagrams, for ease of notational purposes.

Similarly, there is a $\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}$ -Hurewicz map of the form $X \rightarrow \mathrm{TQ}|_{\mathrm{Nil}_{N+1}}(X)$, given by the map $\tilde{X} \rightarrow J_1 \circ_{J_N} (\tilde{X})$ in $\mathrm{Alg}_{\mathcal{O}}$, and iterating this Hurewicz map leads to a cosimplicial resolution of X with respect to $\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}$ -homology of the form

$$X \longrightarrow \mathrm{TQ}|_{\mathrm{Nil}_{N+1}}(X) \rightrightarrows (\mathrm{TQ}|_{\mathrm{Nil}_{N+1}})^2(X) \rightrightarrows (\mathrm{TQ}|_{\mathrm{Nil}_{N+1}})^3(X) \cdots$$

given by the map of cosimplicial J_N -algebras (and by restriction along $\mathcal{O} \rightarrow J_N$, also a map of cosimplicial \mathcal{O} -algebras)

$$\tilde{X} \longrightarrow J_1 \circ_{J_N} (\tilde{X}) \rightrightarrows J_1 \circ_{J_N} J_1 \circ_{J_N} (\tilde{X}) \rightrightarrows \cdots \quad (2.8)$$

where \tilde{X} is regarded as a constant cosimplicial diagram, and applying $\mathrm{holim}_{\Delta}(-)$ to the map of cosimplicial \mathcal{O} -algebras gives the $\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}$ -completion map of the form $X \rightarrow X_{\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}}^{\wedge}$.

The map of operads $\mathcal{O} \rightarrow J_N$ induces a commutative diagram of the form

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & J_1 \circ_{\mathcal{O}} (\tilde{X}) \rightrightarrows J_1 \circ_{\mathcal{O}} J_1 \circ_{\mathcal{O}} (\tilde{X}) \rightrightarrows \cdots \\ \parallel & & \downarrow \qquad \qquad \qquad \downarrow \\ \tilde{X} & \longrightarrow & J_1 \circ_{J_N} (\tilde{X}) \rightrightarrows J_1 \circ_{J_N} J_1 \circ_{J_N} (\tilde{X}) \rightrightarrows \cdots \end{array}$$

and applying $\mathrm{holim}_{\Delta}(-)$ gives the left-hand commutative diagram of the form (2.9).

Theorem 2.12 (Nilpotent TQ-completion retract theorem). Let \mathcal{O} be an operad in \mathcal{R} -modules with $\mathcal{O}[0] = *$. Let $N \geq 1$. If X is an $(N + 1)$ -nilpotent \mathcal{O} -algebra, then there is a left-hand commutative diagram of the form

$$\begin{array}{ccc} X & \longrightarrow & X_{\mathrm{TQ}}^{\wedge} \\ \parallel & & \downarrow \\ X & \xrightarrow[\simeq]{(*)} & X_{\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}}^{\wedge} \end{array} \qquad \begin{array}{ccc} & \text{id} & \\ & \curvearrowright & \\ X & \longrightarrow & X_{\mathrm{TQ}}^{\wedge} \longrightarrow X \end{array} \quad (2.9)$$

in $\mathbf{Alg}_{\mathcal{O}}$ with natural map $(*)$ a weak equivalence. In particular, the TQ-completion map $X \rightarrow X_{\mathbf{TQ}}^{\wedge}$ fits into the right-hand retract diagram in the homotopy category $\mathbf{Ho}(\mathbf{Alg}_{\mathcal{O}})$, natural in X . Here we are assuming that \mathcal{O} satisfies Assumption 2.1.

The purpose of the rest of this section is to prove Theorems 2.12 and 1.1. Let's verify that the comparison map $(*)$ is a weak equivalence. Our approach is to resolve the coaugmented cosimplicial diagram (2.8) of J_N -algebras term-by-term with the finite tower $\{J_n \circ_{J_N} (-)\}_n$, where $1 \leq n \leq N$; we are motivated by the work in [1]. This leads to the commutative diagram in $\mathbf{Alg}_{\mathcal{O}}$ of the form

$$\begin{array}{ccccccc}
\tilde{X} & \xrightarrow{(\#)} & J_1 \circ_{J_N} (\tilde{X}) & \rightrightarrows & J_1 \circ_{J_N} J_1 \circ_{J_N} (\tilde{X}) & \rightleftarrows & \cdots \\
\parallel & & \parallel & & \parallel & & \\
J_N \circ_{J_N} (\tilde{X}) & \xrightarrow{(\#)_N} & J_N \circ_{J_N} (J_1 \circ_{J_N} (\tilde{X})) & \rightrightarrows & J_N \circ_{J_N} (J_1 \circ_{J_N} J_1 \circ_{J_N} (\tilde{X})) & \rightleftarrows & \cdots \\
\downarrow & & \downarrow & & \downarrow & & \\
\vdots & & \vdots & & \vdots & & \cdots \\
\downarrow & & \downarrow & & \downarrow & & \\
J_2 \circ_{J_N} (\tilde{X}) & \xrightarrow{(\#)_2} & J_2 \circ_{J_N} (J_1 \circ_{J_N} (\tilde{X})) & \rightrightarrows & J_2 \circ_{J_N} (J_1 \circ_{J_N} J_1 \circ_{J_N} (\tilde{X})) & \rightleftarrows & \cdots \\
\downarrow & & \downarrow & & \downarrow & & \\
J_1 \circ_{J_N} (\tilde{X}) & \xrightarrow{(\#)_1} & J_1 \circ_{J_N} (J_1 \circ_{J_N} (\tilde{X})) & \rightrightarrows & J_1 \circ_{J_N} (J_1 \circ_{J_N} J_1 \circ_{J_N} (\tilde{X})) & \rightleftarrows & \cdots
\end{array}$$

We will show by induction up the finite tower that $\mathrm{holim}_{\Delta}(\#)_n$ is a weak equivalence for each $1 \leq n \leq N$. Since the bottom horizontal coaugmented cosimplicial diagram has extra codegeneracy maps s^{-1} ([10, 6.2]) induced by the map $J_N \rightarrow J_1$, by cofinality [9, 3.16] it follows that $\mathrm{holim}_{\Delta}(\#)_1$ is a weak equivalence. To make further progress, we must identify the homotopy fibers of these towers.

Proposition 2.13. Let Z be a cofibrant J_N -algebra. There is a zigzag of weak equivalences of towers of the form

$$\{\tau_n J_N \circ_{J_N} (Z)\}_n \simeq \{J_n \circ_{J_N} (Z)\}_n$$

natural with respect to all such Z ; here, $1 \leq n \leq N$.

Proof. There is a zigzag of weak equivalences of the form

$$\begin{array}{c}
\tau_n J_N \circ_{J_N} (Z) \xleftarrow{\simeq} |\mathrm{Bar}(\tau_n J_N, J_N, Z)| \xrightarrow{\simeq} \\
|\mathrm{Bar}(\tau_n \mathcal{O}, J_N, Z)| \xleftarrow{\simeq} |\mathrm{Bar}(J_n, J_N, Z)| \xrightarrow{\simeq} J_n \circ_{J_N} (Z)
\end{array}$$

natural in all such Z ; we are freely using the ‘‘Bar’’ notation from [18, 4.10].

Q.E.D.

Proposition 2.14. Let Z be a cofibrant J_N -algebra. For each $2 \leq n \leq N$, there is a homotopy fiber sequence of the form

$$F_n(Z) \rightarrow J_n \circ_{J_N} (Z) \rightarrow J_{n-1} \circ_{J_N} (Z)$$

in $\mathbf{Alg}_{\mathcal{O}}$, natural in all such Z , where $F_n(Z)$ is defined by

$$F_n(Z) := i_n J_N \circ_{\tau_1 J_N}^h (\tau_1 J_1 \circ_{J_1} J_1 \circ_{J_N} (Z)) \quad (2.10)$$

Proof. By Proposition 2.14, together with [18, 4.21], there is a homotopy fiber sequence of the form

$$i_n J_N \circ_{\tau_1 J_N}^h (\tau_1 J_N \circ_{J_N} (Z)) \rightarrow J_n \circ_{J_N} (Z) \rightarrow J_{n-1} \circ_{J_N} (Z)$$

in $\mathbf{Alg}_{\mathcal{O}}$, natural in all such Z . Noting that the weak equivalence $\tau_1 J_N \xrightarrow{\simeq} \tau_1 J_1$ induces a weak equivalence of the form

$$\begin{aligned} i_n J_N \circ_{\tau_1 J_N}^h (\tau_1 J_N \circ_{J_N} (Z)) &\simeq i_n J_N \circ_{\tau_1 J_N}^h (\tau_1 J_1 \circ_{J_N} (Z)) \\ &\simeq i_n J_N \circ_{\tau_1 J_N}^h (\tau_1 J_1 \circ_{J_1} J_1 \circ_{J_N} (Z)) = F_n(Z) \end{aligned}$$

completes the proof. Q.E.D.

Proof of Theorem 2.12. Consider the commutative diagram of the form

$$\begin{array}{ccccccc} F_n(\tilde{X}) & \xrightarrow{(*)_n} & F_n(J_1 \circ_{J_N}(\tilde{X})) & \rightrightarrows & F_n(J_1 \circ_{J_N} J_1 \circ_{J_N}(\tilde{X})) & \rightrightarrows & \cdots \\ \downarrow & & \downarrow & & \downarrow & & \\ J_n \circ_{J_N}(\tilde{X}) & \xrightarrow{(\#)_n} & J_n \circ_{J_N}(J_1 \circ_{J_N}(\tilde{X})) & \rightrightarrows & J_n \circ_{J_N}(J_1 \circ_{J_N} J_1 \circ_{J_N}(\tilde{X})) & \rightrightarrows & \cdots \\ \downarrow & & \downarrow & & \downarrow & & \\ J_{n-1} \circ_{J_N}(\tilde{X}) & \xrightarrow{(\#)_{n-1}} & J_{n-1} \circ_{J_N}(J_1 \circ_{J_N}(\tilde{X})) & \rightrightarrows & J_{n-1} \circ_{J_N}(J_1 \circ_{J_N} J_1 \circ_{J_N}(\tilde{X})) & \rightrightarrows & \cdots \end{array} \quad (2.11)$$

in $\mathbf{Alg}_{\mathcal{O}}$, for each $(2 \leq n \leq N)$. We know that the coaugmented cosimplicial diagram in the top horizontal row has extra codegeneracy maps s^{-1} ([10, 6.2]), since it has the form of a functor $i_n J_N \circ_{\tau_1 J_N}^h (\tau_1 J_1 \circ_{J_1} -)$ applied to the coaugmented cosimplicial diagram of the form

$$J_1 \circ_{J_N}(\tilde{X}) \xrightarrow{(\#)_1} J_1 \circ_{J_N}(J_1 \circ_{J_N}(\tilde{X})) \rightrightarrows J_1 \circ_{J_N}(J_1 \circ_{J_N} J_1 \circ_{J_N}(\tilde{X})) \rightrightarrows \cdots$$

in \mathbf{Alg}_{J_1} which has extra codegeneracy maps s^{-1} ([10, 6.2]) in \mathbf{Alg}_{J_1} induced by the map $J_N \rightarrow J_1$. It follows from cofinality [9, 3.16] that $\mathrm{holim}_{\Delta}(\#)_1$ and $\mathrm{holim}_{\Delta}(\#)_n$ are weak equivalences for each $2 \leq n \leq N$. Applying holim_{Δ} to (2.11), regarded as a commutative diagram of Δ -shaped diagrams

in $\mathbf{Alg}_{\mathcal{O}}$, we have a commutative diagram of the form

$$\begin{array}{ccc}
F_n(\tilde{X}) & \xrightarrow[\simeq]{\mathrm{holim}_{\Delta}(*)_n} & \mathrm{holim}_{\Delta}\left(F_n(J_1 \circ_{J_N}(\tilde{X})) \Rightarrow \cdots\right) \\
\downarrow & & \downarrow \\
J_n \circ_{J_N}(\tilde{X}) & \xrightarrow{\mathrm{holim}_{\Delta}(\#)_n} & \mathrm{holim}_{\Delta}\left(J_n \circ_{J_N}(J_1 \circ_{J_N}(\tilde{X})) \Rightarrow \cdots\right) \\
\downarrow & & \downarrow \\
J_{n-1} \circ_{J_N}(\tilde{X}) & \xrightarrow{\mathrm{holim}_{\Delta}(\#)_{n-1}} & \mathrm{holim}_{\Delta}\left(J_{n-1} \circ_{J_N}(J_1 \circ_{J_N}(\tilde{X})) \Rightarrow \cdots\right)
\end{array} \tag{2.12}$$

in $\mathbf{Alg}_{\mathcal{O}}$, with vertical columns homotopy fiber sequences, since homotopy limits commute, up to weak equivalence. Consider diagram (2.12) in the case $n = 2$; then since the top and bottom horizontal maps are weak equivalences, it follows that the middle horizontal map $\mathrm{holim}_{\Delta}(\#)_2$ is a weak equivalence; we are using the fact that every homotopy fiber sequence has an associated long exact sequence in (derived or true) homotopy groups of spectra, together with the five lemma from homological algebra (see, for instance, [31, 32]). Hence, by induction up the tower, it follows that $\mathrm{holim}_{\Delta}(\#)_n$ is a weak equivalence for each $1 \leq n \leq N$. In particular, $\mathrm{holim}_{\Delta}(\#) = \mathrm{holim}_{\Delta}(\#)_N$ is a weak equivalence, but this is precisely the map $X \xrightarrow{\simeq} X_{\mathrm{TQ}|_{\mathrm{Nil}_{N+1}}}^{\wedge}$ in diagram (2.9), which completes the proof. Q.E.D.

Proof of Theorem 1.1. The ‘‘only if’’ direction is trivial since TQ-homology preserves weak equivalences by construction. Conversely, suppose f induces a weak equivalence $\mathrm{TQ}(X) \simeq \mathrm{TQ}(Y)$ on TQ-homology. Consider the commutative diagram of the form (by Theorem 2.12)

$$\begin{array}{ccccccc}
X & \longrightarrow & X_{\mathrm{TQ}}^{\wedge} & \longrightarrow & X_{\mathrm{TQ}|_{\mathrm{Nil}_M}}^{\wedge} & \xleftarrow{\simeq} & X \\
f \downarrow & & \simeq \downarrow (\#) & & \downarrow & & \downarrow f \\
Y & \longrightarrow & Y_{\mathrm{TQ}}^{\wedge} & \longrightarrow & Y_{\mathrm{TQ}|_{\mathrm{Nil}_M}}^{\wedge} & \xleftarrow{\simeq} & Y
\end{array} \tag{2.13}$$

Since $\mathrm{TQ}(X) \rightarrow \mathrm{TQ}(Y)$ is a weak equivalence, it follows that $(\#)$ is a weak equivalence, and hence f is a weak equivalence; this is because diagram (2.13) is a retract diagram by (2.9) after passing to the homotopy category $\mathbf{Ho}(\mathbf{Alg}_{\mathcal{O}})$. Q.E.D.

The purpose of the rest of this section is to point out, for completeness, that a similar retract property is also true with TQ-completion replaced by homotopy completion.

Proposition 2.15. Let Z be a J_N -algebra. There are maps of towers of the form

$$\{\tau_n \mathcal{O} \circ_{\mathcal{O}}(Z)\}_n \rightarrow \{\tau_n J_N \circ_{\mathcal{O}}(Z)\}_n \rightarrow \{\tau_n J_N \circ_{J_N}(Z)\}_n$$

natural with respect to Z .

Proof. This is because the left-hand map is induced by $\tau_n \mathcal{O} \rightarrow \tau_n J_N$ and the right-hand map is induced by $\mathcal{O} \rightarrow J_N$. Q.E.D.

It follows that there is a commutative diagram of towers

$$\begin{array}{ccc} \{\tilde{X}\}_n & \longrightarrow & \{\tau_n \mathcal{O} \circ_{\mathcal{O}} (\tilde{X})\}_n \\ \parallel & & \downarrow \\ \{\tilde{X}\}_n & \longrightarrow & \{\tau_n J_N \circ_{J_N} (\tilde{X})\}_n \end{array}$$

of \mathcal{O} -algebras, where $\{\tilde{X}\}_n$ denotes the constant tower with value \tilde{X} , and applying holim_n gives the left-hand commutative diagram of the form (2.14).

Theorem 2.16 (Nilpotent homotopy completion retract theorem). Let \mathcal{O} be an operad in \mathcal{R} -modules with $\mathcal{O}[0] = *$. Let $N \geq 1$. If X is an $(N + 1)$ -nilpotent \mathcal{O} -algebra, then there is a left-hand commutative diagram of the form

$$\begin{array}{ccc} X & \longrightarrow & X^{\text{h}\wedge} \\ \parallel & & \downarrow \\ X & \xrightarrow[\simeq]{(*)} & X^{\text{h}\wedge}|_{\text{Nil}_{N+1}} \end{array} \qquad \begin{array}{ccc} & \text{id} & \\ & \curvearrowright & \\ X & \longrightarrow & X^{\text{h}\wedge} \longrightarrow X \end{array} \tag{2.14}$$

in $\text{Alg}_{\mathcal{O}}$ with natural map $(*)$ a weak equivalence. In particular, the homotopy completion map $X \rightarrow X^{\text{h}\wedge}$ fits into the right-hand retract diagram in the homotopy category $\text{Ho}(\text{Alg}_{\mathcal{O}})$, natural in X . Here we are assuming that \mathcal{O} satisfies Assumption 2.1.

Proof. It suffices to observe that $(*)$ is a weak equivalence. This follows from the fact that $\tilde{X} \xrightarrow{\simeq} \tau_n J_N \circ_{J_N} (\tilde{X})$ is a weak equivalence for each $n \geq N$; this follows from the observation that the map $J_N \xrightarrow{\simeq} \tau_n J_N$ is a weak equivalence for each $n \geq N$. Q.E.D.

References

- [1] G. Arone and M. Kankaanrinta. A functorial model for iterated Snaith splitting with applications to calculus of functors. In *Stable and unstable homotopy (Toronto, ON, 1996)*, volume 19 of *Fields Inst. Commun.*, pages 1–30. Amer. Math. Soc., Providence, RI, 1998.
- [2] M. Basterra. André-Quillen cohomology of commutative S -algebras. *J. Pure Appl. Algebra*, 144(2):111–143, 1999.
- [3] M. Basterra and M. A. Mandell. Homology and cohomology of E_{∞} ring spectra. *Math. Z.*, 249(4):903–944, 2005.
- [4] M. Basterra and M. A. Mandell. Homology of E_n ring spectra and iterated THH . *Algebr. Geom. Topol.*, 11(2):939–981, 2011.
- [5] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin, 1972.

- [6] M. Ching and J. E. Harper. Derived Koszul duality and TQ-homology completion of structured ring spectra. [arXiv:1502.06944 \[math.AT\]](#), 2015.
- [7] M. Ching and J. E. Harper. Higher homotopy excision and Blakers-Massey theorems for structured ring spectra. *Adv. Math.*, 298:654–692, 2016.
- [8] E. Dror. A generalization of the Whitehead theorem. In *Symposium on Algebraic Topology (Battelle Seattle Res. Center, Seattle, Wash., 1971)*, pages 13–22. Lecture Notes in Math., Vol. 249. Springer, Berlin, 1971.
- [9] E. Dror and W. G. Dwyer. A long homology localization tower. *Comment. Math. Helv.*, 52(2):185–210, 1977.
- [10] W. G. Dwyer, H. R. Miller, and J. Neisendorfer. Fibrewise completion and unstable Adams spectral sequences. *Israel J. Math.*, 66(1-3):160–178, 1989.
- [11] B. Fresse. Lie theory of formal groups over an operad. *J. Algebra*, 202(2):455–511, 1998.
- [12] B. Fresse. Koszul duality of operads and homology of partition posets. In *Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory*, volume 346 of *Contemp. Math.*, pages 115–215. Amer. Math. Soc., Providence, RI, 2004.
- [13] P. G. Goerss. On the André-Quillen cohomology of commutative \mathbf{F}_2 -algebras. *Astérisque*, (186):169, 1990.
- [14] T. G. Goodwillie. Calculus. III. Taylor series. *Geom. Topol.*, 7:645–711, 2003.
- [15] J. E. Harper. Homotopy theory of modules over operads in symmetric spectra. *Algebr. Geom. Topol.*, 9(3):1637–1680, 2009. Corrigendum: *Algebr. Geom. Topol.*, 15(2):1229–1237, 2015.
- [16] J. E. Harper. Bar constructions and Quillen homology of modules over operads. *Algebr. Geom. Topol.*, 10(1):87–136, 2010.
- [17] J. E. Harper. Homotopy theory of modules over operads and non- Σ operads in monoidal model categories. *J. Pure Appl. Algebra*, 214(8):1407–1434, 2010.
- [18] J. E. Harper and K. Hess. Homotopy completion and topological Quillen homology of structured ring spectra. *Geom. Topol.*, 17(3):1325–1416, 2013.
- [19] M. Hovey, B. Shipley, and J. H. Smith. Symmetric spectra. *J. Amer. Math. Soc.*, 13(1):149–208, 2000.
- [20] N. J. Kuhn. Localization of André-Quillen-Goodwillie towers, and the periodic homology of infinite loopspaces. *Adv. Math.*, 201(2):318–378, 2006.
- [21] N. J. Kuhn. Goodwillie towers and chromatic homotopy: an overview. In *Proceedings of the Nishida Fest (Kinosaki 2003)*, volume 10 of *Geom. Topol. Monogr.*, pages 245–279. Geom. Topol. Publ., Coventry, 2007.
- [22] N. J. Kuhn. Adams filtration and generalized Hurewicz maps for infinite loopspaces. [arXiv:1403.7501 \[math.AT\]](#), 2014.

- [23] N. J. Kuhn and L. A. Pereira. Operad bimodules and composition products on André-Quillen filtrations of algebras. *Algebr. Geom. Topol.*, 17(2):1105–1130, 2017.
- [24] R. McCarthy and V. Minasian. On triples, operads, and generalized homogeneous functors. [arXiv:math/0401346v1](https://arxiv.org/abs/math/0401346v1) [math.AT], 2004.
- [25] H. R. Miller. The Sullivan conjecture on maps from classifying spaces. *Ann. of Math. (2)*, 120(1):39–87, 1984. Correction: *Ann. of Math. (2)*, 121(3):605–609, 1985.
- [26] D. Pavlov and J. Scholbach. Homotopy theory of symmetric powers. *Homology Homotopy Appl.*, 20(1):359–397, 2018.
- [27] L. A. Pereira. Goodwillie calculus in the category of algebras over a spectral operad. 2013. Available at: <http://math.mit.edu/~luisalex/>.
- [28] L. A. Pereira. Cofibrancy of operadic constructions in positive symmetric spectra. *Homology Homotopy Appl.*, 18(2):133–168, 2016.
- [29] D. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [30] C. Rezk. *Spaces of Algebra Structures and Cohomology of Operads*. PhD thesis, MIT, 1996. Available at <http://www.math.uiuc.edu/~rezk/>.
- [31] S. Schwede. *An untitled book project about symmetric spectra*. 2007,2009. Available at: <http://www.math.uni-bonn.de/people/schwede/>.
- [32] S. Schwede. On the homotopy groups of symmetric spectra. *Geom. Topol.*, 12(3):1313–1344, 2008.
- [33] B. Shipley. A convenient model category for commutative ring spectra. In *Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory*, volume 346 of *Contemp. Math.*, pages 473–483. Amer. Math. Soc., Providence, RI, 2004.